

Automorphisms of seamed surfaces, modular operads and Galois actions

Marcy Robertson (University of Melbourne)

joint with: Boavida, Bonatto, Hackney, Horel and Yau

A really fast introduction to a lot of cool math:

- Let $Gal(\mathbb{Q})$ denote the absolute Galois group of \mathbb{Q} .
- This is a large profinite group:

$$\widehat{G} = \varprojlim G/H$$

but we don't even know the finite quotients of $Gal(\mathbb{Q})$!

Idea : Identify $g \in Gal(\mathbb{Q})$ with a pair

$$(\chi(g), f_g) \in \widehat{\mathbb{Z}}^* \times \widehat{F}'_2$$

- $\chi(g)$ is the **cyclotomic character**.
- $\widehat{F}'_2 = \pi_1(\mathcal{M}_{0,4}) \cong \widehat{\Gamma}_{0,4}$.

A slightly easier group: \widehat{GT}

Notation: For any homomorphism of profinite groups

$$\widehat{F}_2 \longrightarrow G$$

$$(x, y) \longmapsto (a, b)$$

we write $f(a, b)$ for the image of any $f \in \widehat{F}_2$. For example:

- Given $id : \widehat{F}_2 \rightarrow \widehat{F}_2$, we have $f = f(x, y)$;
- Given the map $\widehat{F}_2 \rightarrow \widehat{F}_2$ which swaps generators x and y we have $f \mapsto f(y, x)$.

A slightly easier group: \widehat{GT}

The **Grothendieck-Teichmüller group** \widehat{GT} is the group of pairs

$$(\lambda, f) \in \widehat{\mathbb{Z}}^* \times \widehat{F}'_2$$

satisfying the property that

$$x \mapsto x^\lambda \quad \text{and} \quad y \mapsto f^{-1}y^\lambda f$$

induce an automorphism of \widehat{F}_2 and :

- (I) $f(x, y)f(y, x) = 1$,
- (II) $f(x, y)x^m f(z, x)z^m f(y, z)y^m = 1$ where $xyz = 1$ and $m = (\lambda - 1)/2$,
- (III) $f(x_{34}, x_{45})f(x_{51}, x_{12})f(x_{23}, x_{34})f(x_{45}, x_{51})f(x_{12}, x_{23}) = 1$ in $\Gamma_{0,5}$
where x_{ij} is a Dehn twist of boundaries i and j .


Theorem (Ihara)

There is an injection $\text{Gal}(\mathbb{Q}) \hookrightarrow \widehat{GT}$.

So our question becomes: What is \widehat{GT} ?


- $\widehat{F}_2 \cong \widehat{\Gamma}_{0,4}$
- relations in \widehat{GT} are coming from mapping class groups.
- The mapping class group has a presentation

$$\Gamma_{g,n} = \langle \alpha_1, \dots, \alpha_k \mid (C), (B), (D), (L) \rangle.$$




Pants Decompositions

A **pants decomposition** of $\Sigma_{g,n}$ is a collection of simple closed curves that cuts $\Sigma_{g,n}$ into *pairs of pants* (i.e. $\Sigma_{0,3}$).




We break Mapping Class Groups into Pants Decompositions

Notice that a **pants decomposition** looks like the result of a composition.



We break Mapping Class Groups into Pants Decompositions

Notice that a **pants decomposition** looks like the result of a composition.




Modular Operads

A \mathfrak{C} -coloured **cyclic operad** is an algebraic structure consisting of:

- an involutive set of colours \mathfrak{C} ;
- for each $c_1, \dots, c_n \in \mathfrak{C}$ a Σ_n -set $P(c_1, \dots, c_n)$;
- a family of equivariant, associative and unital composition operations

$$P(c_1, \dots, c_n) \times P(d_1, \dots, d_m) \longrightarrow P(c_1, \dots, \hat{c}_i, \dots, d_1, \dots, \hat{d}_j, \dots, d_m),$$

when $c_i = d_j^\dagger$.




A \mathcal{C} -coloured **modular operad** is a cyclic operad which also has

- a family of equivariant contraction operations

$$P(c_1, \dots, c_n) \longrightarrow P(c_1, \dots, \hat{c}_i, \dots, \hat{c}_j, \dots, c_n),$$

when $c_i = c_j^\dagger$.


satisfying some axioms.



A modular operad is almost the right thing for studying $\widehat{\Gamma}_{g,n}$

There is an equivalence of categories:

$$\text{ModOp} \xrightarrow[\cong]{N} \mathbf{Set}_{\text{Segal}}^{\text{U}^{\text{op}}}$$




Graphs:


A **graph** G is a diagram of finite sets:

$$i \curvearrowright A \xleftarrow{s} D \xrightarrow{t} V$$

- i is a free involution;
- s is a monomorphism.




Graphical Maps



Inner coface maps


An **inner coface map** $d_v : G \rightarrow G'$ is a graphical map defined by “blowing-up” a single vertex v in G by a graph which has precisely **one** internal edge.



Outer coface maps

An **outer coface map** is either:

- an **embedding** $d_e : G \rightarrow G'$ in which G' has precisely **one** more internal edge than G or
- an **embedding** $\downarrow \rightarrow \star_n$.



Codegeneracy maps

A **codegeneracy map** $s_v : G \rightarrow G'$ is a graphical map defined by “blowing-up” a vertex v in G by \downarrow .




The graphical category

The **graphical category** \mathcal{U} is the category whose objects are connected graphs. The morphisms are composites of inner coface maps, outer coface maps, codegeneracies and isomorphisms.


Graphical Sets




The category of **graphical sets** is $\text{Set}^{\mathcal{U}^{op}}$.

- X_G : evaluation of X at $G \in \mathcal{U}$.
- $\varphi : G \rightarrow G' \Rightarrow \varphi^* : X_{G'} \rightarrow X_G$.




A special graphical set made of internal edges: $X_G^1 = \lim_{\star_v \leftarrow \updownarrow \rightarrow \star_w}$





Segal Maps



The embeddings $\star_v \hookrightarrow G$ induce a Segal map

$$X_G \longrightarrow X_G^1 \subseteq \prod_{v \in V(G)} X_{\star_v}$$

which factors through X_G^1 .

Segal graphical sets

A graphical set $X \in \mathbf{Set}^{\mathbf{U}^{op}}$ is strictly **Segal** if the Segal map

$$X_G \longrightarrow X_G^1 \subseteq \prod_{v \in V(G)} X_{\star_v}$$

is a bijection for each G in \mathbf{U} .

Theorem (HRY20b)

There is an equivalence of categories:

$$\mathbf{ModOp} \xrightarrow[\cong]{N} \mathbf{Set}_{\mathbf{Segal}}^{\mathbf{U}^{op}}.$$

A graphical set $X \in \mathbf{sSet}^{\mathbf{U}^{op}}$ is weakly **Segal** if the Segal map

$$X_G \longrightarrow X_G^1 \subseteq \prod_{v \in V(G)} X_{\star_v}$$

is a weak homotopy equivalence for each G in \mathbf{U} .

Weak Segal Modular Operads Take Us Back To $Gal(\mathbb{Q})$

A modular operad of Seamed Surfaces

The goupoid $\mathcal{S}_{g,n}$:

- objects are surfaces $P := (\Sigma_{g,n}, P, Q)$ together with a “atomic” **pants decomposition**;
- morphisms are $\pi_0 \text{Diff}^+(\Sigma_{g,n}, \partial, \sigma)$.

Σ_n acts freely on $\mathcal{S}_{g,n}$ by permuting the labels of boundaries \Rightarrow

$$B\mathcal{S}_{g,n} \simeq B\Gamma_{g,n}.$$

A modular operad of Seamed Surfaces

Operations:

$$\mathcal{S}_{g,n} \times_{ij} \mathcal{S}_{h,k} \xrightarrow{\circ_{ij}} \mathcal{S}_{g+h,n+k-2}$$

and

$$\mathcal{S}_{g,n} \xrightarrow{\xi_{ij}} \mathcal{S}_{g+1,n-2}$$

can be defined on objects by gluing surfaces and on morphisms as the "combination" of the maps on the subsurfaces.

These are well-defined, associative operations and thus

$$\mathcal{S} = \{\mathcal{S}_{g,n}\}$$

assembles into a **modular operad in groupoids**. \Rightarrow

$$NS = \{NS_{g,n}\} \in \text{Set}_{\text{Segal}}^{\text{U}^{\text{op}}}$$

Proposition (BHR)

$$\widehat{GT} \cong \pi_0 \text{map}^h(\widehat{NS}_0, \widehat{NS}_0)$$

But to get back to our comparison with the mapping class groups:

Theorem (BHR)

$$\widehat{GT} \cong \pi_0 \text{map}^h(B\widehat{NS}_0, B\widehat{NS}_0)$$

Proposition (BHR)

$$\widehat{GT} \cong \pi_0 \operatorname{map}^h(\widehat{NS}_0, \widehat{NS}_0)$$

But to get back to our comparison with the mapping class groups:

Theorem (BHR)

$$\widehat{GT} \cong \pi_0 \operatorname{map}^h(B\widehat{NS}_0, B\widehat{NS}_0)$$

Point: Here we can see how $\operatorname{Gal}(\mathbb{Q})$ acts.

Groups Related to \widehat{GT} : The higher genus case

There is a subgroup $\Lambda \subseteq \widehat{GT}$:

Theorem (BR)

There is an isomorphism

$$\Lambda \cong \text{End}_0(N\widehat{S}).$$

Theorem (BR - In Progress)

There is an isomorphism

$$\Lambda \cong \pi_0 \text{map}^h(B\widehat{NS}).$$

Question: Can weak Segal modular operads give us even more information about $\text{Gal}(\mathbb{Q})$?

Thanks!